



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Sex differences in leukocyte telomere length in a free-living mammal

**Citation for published version:**

Watson, RL, Bird, EJ, Underwood, S, Adams, RV, Fairlie, J, Watt, K, Salvo-Chirnside, E, Pilkington, J, Pemberton, J, McNeilly, TN, Froy, H & Nussey, D 2017, 'Sex differences in leukocyte telomere length in a free-living mammal', *Molecular Ecology*. <https://doi.org/10.1111/mec.13992>

**Digital Object Identifier (DOI):**

[10.1111/mec.13992](https://doi.org/10.1111/mec.13992)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Molecular Ecology

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Sex differences in leukocyte telomere length in a free-living mammal

Watson, R.L.<sup>1</sup>, Bird, E.J.<sup>1</sup>, Underwood, S.<sup>1</sup>, Adams, R.V.<sup>1</sup>, Fairlie, J.<sup>1</sup>, Watt, K.<sup>1</sup>, Salvo-Chirnside, E.<sup>2</sup>, Pilkington, J.G.<sup>1</sup>, Pemberton, J.M.<sup>1</sup>, McNeilly, T.N.<sup>3</sup>, Froy, H.<sup>1</sup> & Nussey, D.H.<sup>1</sup>

<sup>1</sup> Institute of Evolutionary Biology, University of Edinburgh, The King's Buildings, Ashworth Laboratories, Charlotte Auerbach Road, Edinburgh EH9 3FL, U.K.

<sup>2</sup> SynthSys, University of Edinburgh, The King's Buildings, Waddington Building, Max Bourne Crescent, Edinburgh EH9 3BF, U.K.

<sup>3</sup> Moredun Research Institute, Pentlands Science Park, Bush Loan, Midlothian EH26 0PZ, U.K.

**Key words:** Lymphocyte, Granulocyte, Soay sheep, Sexual selection, polygyny, *Ovis aries*

**Corresponding Author:** Daniel Nussey, University of Edinburgh, The King's Buildings, Ashworth Laboratories, Charlotte Auerbach Road, Edinburgh EH9 3FL. Tel: +44 (0)131 6505494. Email: [dan.nussey@ed.ac.uk](mailto:dan.nussey@ed.ac.uk)

**Running Title:** Sex differences in telomere length

## Abstract

Mounting evidence suggests that average telomere length reflects previous stress and predicts subsequent survival across vertebrate species. In humans, leukocyte telomere length (LTL) is consistently shorter during adulthood in males than females, although the causes of this sex difference and its generality to other mammals remain unknown. Here we measured LTL in a cross sectional sample of free-living Soay sheep and found shorter telomeres in males than females in later adulthood (>3 years of age), but not in early life. This observation was not related to sex differences in growth or parasite burden, but we did find evidence for reduced LTL associated with increased horn growth in early life in males. Variation in LTL was independent of variation in the proportions of different leukocyte cell types, which are known to differ in telomere length. Our results provide the first evidence of sex differences in LTL from a wild mammal, but longitudinal studies are now required to determine whether telomere attrition rates or selective disappearance are responsible for these observed differences.

## Introduction

Telomeres are G-rich repeat sequences of DNA at the ends of chromosomes, which form complexes with proteins that protect linear chromosomes from DNA repair machinery and the so-called 'end replication problem' (Blackburn 1991; De Lange 2004). Telomeres shorten with each cell replication and are also highly sensitive to damage by reactive oxygen species (von Zglinicki 2002), but can be replenished through the action of the enzyme telomerase (Armanios & Blackburn 2012). When telomere length drops below a critical threshold it triggers the onset of cellular senescence, and thus there may be close links between telomere attrition and cellular ageing (Gomes *et al.* 2011; Hemann *et al.* 2001). Average telomere length (TL), which is typically measured from DNA obtained from whole blood samples, has been shown to decline with age in numerous species and is a biomarker of considerable current interest in human epidemiology and evolutionary ecology (Blackburn *et al.* 2015; Monaghan & Haussmann 2006). In humans, TL in adulthood appears predictive of late-onset diseases and, in some studies, mortality risk (Blackburn *et al.* 2015; Cawthon *et al.* 2003; Rode *et al.* 2015), whilst prior experience of chronic stress is associated with shorter TL (Monaghan 2014; Shalev 2012). Studies of wild birds and mammals have also recently found associations between short TL and reduced subsequent survival or recapture rates (Barrett *et al.* 2013; Bize *et al.* 2009; Fairlie *et al.* 2016).

In humans and laboratory rodents, females generally have longer TL than males in adulthood (Barrett & Richardson 2011; Gardner *et al.* 2014). This pattern mirrors the pervasive gender difference in longevity observed in humans and many other mammals, leading to speculation that sex differences in TL may be related to differences in lifespan (Aviv *et al.* 2005; Stindl 2004). A recent meta-analysis

of forty adult human data sets concluded that females have significantly longer telomeres than males (Gardner *et al.* 2014), and more recent large-scale studies have confirmed this pattern (Berglund *et al.* 2016; Lapham *et al.* 2015). However, it remains unclear whether sex differences in TL are present in early life or emerge in later adulthood due to differences in telomere attrition rates in humans. Several studies have failed to document significant sex differences in TL measured in umbilical cord blood (Akkad *et al.* 2006; Aubert *et al.* 2012; Okuda *et al.* 2002; Shi-Ni *et al.* 2013), although a recent larger-scale study did find significantly longer telomeres in new born females than males (Factor-Litvak *et al.* 2016). Studies of captive primates have yielded equivocal results: no sex differences in TL were found in cynomolgus macaques (*Macaca fascicularis*; Gardner *et al.* 2007), whilst female rhesus macaques (*Macaca mulatta*) had significantly longer TL than males in leukocytes but not in other tissue types (Smith *et al.* 2011). In laboratory populations of rats (*Rattus rattus*; Cherif *et al.* 2003; Tarry-Adkins *et al.* 2006) and Algerian mice (*Mus spretus*; Coviello-McLaughlin & Prowse 1997) females were also reported to have longer telomeres than males.. To date, studies of sex differences in TL in non-human mammals remain limited to primates and rodents in captivity: we could find only one study comparing TL between the sexes in a wild mammal population. This study of European badgers, which show strong sexual size dimorphism, found no evidence of sex differences in mean TL or changes in TL with age (*Meles meles*; Beirne *et al.* 2014).

The pattern of sex differences in TL across vertebrate species appears variable, and several non-mutually exclusive explanations have been proposed for this variation, including heterogametic disadvantage and differences in the effects of sex hormones on oxidative stress and telomerase function (Barrett & Richardson 2011; Gardner *et al.* 2014). It has also been hypothesized that, in species with sexual size dimorphism, increased cell proliferation rates and oxidative stress associated with increased growth in the larger sex could reduce TL (Barrett & Richardson 2011; Stindl 2004).

Furthermore, increased parasite burdens in males relative to females have been documented in polygynous mammals and proposed as a driver of male-biased mortality (Moore & Wilson 2002). Experiments in laboratory-kept house mice (*Mus musculus*) demonstrated faster leukocyte TL loss in males than females following repeated bacterial infection, raising the possibility that sex differences in TL could reflect sex differences in the response to infection (Ilmonen *et al.* 2008). Polygynous males also typically exhibit secondary sexual characteristics other than increased body size (e.g. seasonal colouration, vocalisations, ornaments and weapons) and highly energetically expensive intra-sexual competition for access to mates (Andersson 1994). Male investment in these reproductive traits may be associated with increased cellular proliferation or oxidative stress, resulting in faster TL attrition in males relative to females (Barrett & Richardson 2011; Monaghan 2010). A recent review found little consistent evidence that sex differences in TL were associated with heterogamety, the degree of body size dimorphism, or mating system (Barrett & Richardson 2011). However, direct tests for sex-specific associations between TL and either weight, reproductive investment or parasite burden within polygynous species remain rare (Beirne *et al.* 2014; Olsson *et al.* 2011).

In mammals, which have enucleated red blood cells, TL measured in DNA extracted from blood samples reflects the average leukocyte telomere length (LTL). This is in contrast to other vertebrate groups with nucleated erythrocytes, where it is erythrocyte telomere length (ETL) that is predominantly measured in blood. This represents a challenge for mammalian studies, as LTL encompasses a range of different white blood cell types which have different functions and roles within the immune system and show differences in proliferative capacity, telomerase expression and ultimately telomere length (Weng 2001). For instance, in humans and baboons granulocytes have longer telomeres than lymphocytes in adulthood, most likely due to the fact that granulocytes are

terminally differentiated cells whilst lymphocytes have the capacity to rapidly replicate and differentiate (Aubert *et al.* 2012; Baerlocher *et al.* 2007; Kimura *et al.* 2010). Lymphocytes also vary in telomere length, with naïve T cells having longer telomere lengths in comparison to memory T cells, again due to greater proliferative history of the latter (Aubert *et al.* 2012; Weng 2001). The composition of circulating leukocyte cell types can change profoundly with age and vary between sexes (Giefing-Kröll *et al.* 2015; Linton & Dorshkind 2004; Pawelec *et al.* 2010), and changes found in average LTL in relation to age and sex could therefore reflect changes in underlying cell population structure (Weng 2001). However, studies in humans and primates have reported very strong within-individual correlations in TL measured in different leukocyte sub-populations and among different tissues (Aubert *et al.* 2012; Baerlocher *et al.* 2007; Daniali *et al.* 2013; Gardner *et al.* 2007; Kimura *et al.* 2010). Based on this apparent ‘synchrony’ in TL across tissue and cell types, it has been argued that among-individual variation in LTL reflects differences in the TL of the haematopoietic stem cell pool, which is primarily determined genetically and by early life environment (Daniali *et al.* 2013). However, studies investigating the dependence of LTL and its associations with age, sex and other traits on variation in leukocyte population structure in non-primate mammals are currently lacking.

Here, we tested sex differences in LTL in a free-living population of Soay sheep on the St Kilda archipelago. Soay sheep have a polygynous breeding system: males compete for mating opportunities with oestrous females during the autumn rut and have highly skewed reproductive success, with a handful of males obtaining the majority of paternities each year (Clutton-Brock & Pemberton 2004). As is typical in polygynous systems, males are larger and shorter lived than females: by five years of age, males average around 38 kg and females around 24 kg in summer with maximum recorded lifespan of ten years in males and sixteen years in females (Clutton-Brock & Pemberton 2004). This population is parasitized by a variety of Strongyle gastrointestinal nematodes,

and parasite burdens – as estimated by faecal egg counts (FEC) – are predictive of over-winter survival and greater in males than females at all ages (Hayward *et al.* 2009; Wilson *et al.* 2004). Horns, which are grown incrementally each year, are an important secondary sexual trait in males in Soay sheep and horn length is positively associated with subsequent male annual reproductive success (Johnston *et al.* 2013). Both sexes exhibit normal and greatly reduced (scurred or polled) horn growth phenotypes, and the genes underpinning this polymorphism have recently been identified (Johnston *et al.* 2013). We have recently shown that LTL is positively associated with survival in early life in female Soay sheep (Fairlie *et al.* 2016), and that age-related variation in the proportion of different T cell sub-types is present in this population (Nussey *et al.* 2012). However, we have yet to test for sex differences in LTL or examine whether and how LTL is associated with variation in leukocyte cell structure in the population. Here, we examine whether sex differences in LTL are present and whether evident sex differences are dependent on age, weight or Strongyle FEC, as well as testing the hypothesis that costly investment in male horn growth should be reflected by reduced LTL. We also measure the proportions of different leukocyte cell types which are known to differ in telomere dynamics and test whether variation in leukocyte population structure can explain observed associations among LTL, sex, age and investment in secondary sexual characteristics.

## Materials & Methods

### Study system & sample preparation

The Soay sheep is a primitive breed of domestic sheep that has been living on the remote St Kilda archipelago with minimal human management for the last few millennia. Since 1985, the sheep living in the Village Bay area of the main island in the archipelago, Hirta, have been the subject of



individual-based study (Clutton-Brock & Pemberton 2004). As part of this study, individuals are caught and tagged within a few days of birth in spring. An individual's horn type is identified at capture as follows: "normal" horns are sturdy and consist of a bony core covered in a keratin sheath, whilst "scurred" horns consist of keratin but lack a bony core. A "polled" phenotype is present only in females and involves a complete absence of visible horn growth. Around 85% of males and 35% of females in the population have normal horns (Clutton-Brock & Pemberton 2004). Each August, 50-60% of the resident population are caught in temporary corral traps. At capture, blood and faecal samples are taken and each animal is weighed to the nearest 0.1 kg and horn length is measured from the base of the horn, along the outer curvature to the tip, with each annual growth increment noted (Clutton-Brock & Pemberton 2004). This study uses data and samples from animals caught during the Augusts of 2014 (78 males and 174 females) and 2015 (66 males and 174 females).

Two 9ml Lithium Heparin Vacuettes of blood were taken from each individual and kept in a cool box or fridge from the point of sampling until further processing within 24 hours of sampling. The first Vacuette of blood was spun at 1,008 x g for 10 minutes and the plasma layer was then drawn off and replaced by the same quantity of 0.9 % NaCl solution and spun again at 1,008 x g for 10 minutes. The intermediate buffy coat layer, comprising mainly white blood cells, was then drawn off into a 1.5 ml Eppendorf tube and stored at -20 °C until used to assay leukocyte telomere length. Faecal samples were available for 427 of the captured individuals, and Strongyle and Strongyloides FEC were estimated in these samples using a modified McMaster technique (following Gulland & Fox 1992).

#### Leukocyte cell measurements

Within 12 hours of collection, 5 µl of whole blood from the second Vacuette was applied to one end of a standard glass microscope slide. The drop of blood was then spread with the edge of a second

slide at a 45° angle to produce an even film. Slides were air dried overnight and stained using a Quick-Diff Kit stain (Gentaur, London) the following day, as per manufacturer's instructions. Differential white blood cell counts were conducted back in the laboratory in Edinburgh. Briefly, 100 cells were counted at 40x magnification using the "battlement track" method and based on staining and morphology, identified as either lymphocytes, eosinophils or neutrophils (Bain 2014). Basophils and monocytes were observed too rarely to analyse. From this data, we calculated the granulocyte (neutrophils and eosinophils) to lymphocyte ratio (GLR) and this ratio was used in subsequent analyses. Only slides with a clear regular monolayer of cells were counted, and slides with uneven cell density or unclear staining were omitted, leaving 465 GLR measurements available for subsequent analyses. See Watson *et al.* (2016) for information on repeatability of these measurements.

A further 1ml of whole blood from the second Vacuette was used to prepare a formalin fixed sample of lymphocytes, which was stored at 4°C until subsequent flow cytometry analysis back in Edinburgh (following Nussey *et al.* 2012). We were able to estimate the proportions of lymphocytes that were helper T cells (CD4+), cytotoxic T cells (CD8+) and, within each of these cell types the proportion that were putatively naïve cells (CD45RA+). Full methodological details are presented in the on-line supplementary material. From this data we calculated the CD4 to CD8 T cell ratio as well as the proportions of helper T cells and cytotoxic cells that were naïve, and we used these three measurements in subsequent analyses.

#### Telomere length measurement

Genomic DNA was extracted from buffy coat using the Qiagen DNeasy Blood and Tissue Kit following manufacturer's guidelines for animal blood (Cat# 69581, Manchester, UK). The protocol was modified slightly to facilitate sample flow through the spin columns which subsequently improved DNA yield

and purity (see on-line supplementary material). Following DNA extraction and elution in buffer AE (10 mM TrisCl, 0.5 EDTA, pH 9.0), a strict quality control protocol was implemented to determine DNA quality and integrity. First, each sample was individually tested for DNA yield and purity using a Nanodrop ND-1000 9 spectrophotometer (Thermo Scientific, Wilmington DE, USA). Samples yielding < 20 ng/μl were immediately rejected. Samples yielding ≥ 20 ng/μl were checked for DNA purity; acceptable ranges for absorption were 1.7 - 2.0 for 260/280 nm ratio and 1.8 - 2.2 for 260/230 nm ratio. Acceptable samples were then diluted to 10 ng/μl and their DNA integrity assessed by running 20 μl (200 ng total DNA) on a 0.5 % agarose gel. Samples were scored for integrity on a scale of 1 to 5 by visual examination of their DNA crowns, with samples scoring higher than 2 being excluded from further analyses (see Seeker *et al.* 2016 for details). Samples which failed one or more of the above QC measures were re-extracted and if they failed QC a second time they were excluded from the study.

Relative leukocyte telomere length (RTL) was measured using real-time quantitative PCR (qPCR; Cawthon 2002), using protocols we have previously developed and validated in sheep and cattle blood samples (Fairlie *et al.* 2016; Seeker *et al.* 2016). The qPCR method estimates the total amount of telomeric sequence present in a sample relative to the amount of a non-variable copy number reference gene. Note that this method measures both terminal and interstitial telomere sequence, and if interstitial telomeric DNA content varies among or within individuals this could influence our results. In this study we used the beta-2-microglobulin (B2M) as our reference gene, and used primers supplied by Primer Design (Catalogue number: HK-SY-Sh-900, Southampton, UK). For telomeric amplification tel1b (5'-CGG TTT GTT TGG GTT TGG GTT TGG GTT TGG GTT TGG GTT-3') and tel 2b (5'-GGC TTG CCT TAC CCT TAC CCT TAC CCT TAC CCT TAC CCT-3') primers were used Epel *et al.* 2004. Telomere primers were manufactured, HPLC purified and supplied by Integrated DNA

Technologies (IDT, Glasgow, UK). Telomere and reference gene reactions were run in separate wells of the same qPCR plate at a concentration of 300 nM and 900 nM respectively. Samples were diluted to 1 ng/μl with buffer AE just prior to qPCR analysis. Each reaction was prepared using 5 μl of LightCycler 480SYBR Green I Master Mix (Cat # 04887352001, Roche, West Sussex, UK) and 1ng of sample DNA in a total reaction volume of 10 μl. We used 384 well plates which were loaded with sample DNA and master mix using an automated liquid handling robot (Freedom Evo-2 150; Tecan).

Each plate included two calibrator samples (1 ng/μl) to account for plate to plate variation and a non-template control (NTC) consisting of nuclease free water. The calibrator sample was extracted from a large quantity of buffy coat prepared from blood supplied from a single domestic sheep (Cat# SHP-BUFCT-LIHP, Sera Laboratories International LTD, West Sussex, UK). We carried out a large number of extractions from this sample, applied the same quality control as above and then pooled the extracts and aliquoted them for subsequent use. Samples, calibrators and NTCs were all run in triplicate. All qPCRs were performed using a Roche LC480 instrument using the following reaction protocol; 10 min at 95 °C (enzyme activation), followed by 50 cycles of 15 s at 95 °C (denaturation) and 30 s at 58 °C (primer annealing), then 30 s at 72 °C (signal acquisition). Melting curve protocol was 1 min at 95 °C, followed by 30 s at 58 °C, then 0.11 °C/s to 95 °C followed by 10 s at 40 °C.

We used the LinRegPCR software package (version 2016.0; Ruijter *et al.* 2009) to correct our amplification curves for baseline fluorescence, and to calculate well-specific reaction efficiencies and Cq values. A constant fluorescence threshold was set within the window of linearity for each amplicon group, calculated using the average Cq across the first 6 plates. The threshold values used were 0.193 and 0.222, and the average efficiency across all plates were 1.88 and 1.91 for the B2M and telomere amplicon groups, respectively. Samples were excluded from further analysis if the

coefficient of variation (CV) across triplicate Cq values for either amplicon was > 5 %, or if at least one of their triplicate reactions had an efficiency that was 5 % higher or lower than the mean efficiency across all wells on that plate for the respective amplicon. Overall, nine samples were excluded based on quality control failure at either extraction or qPCR stages, leaving 492 samples available for use in further analyses.

Relative LTL for each sample was calculated, following Pfaffl 2001, using average reaction efficiencies for each plate and Cq for each sample determined by LinRegPCR as follows:

$$RTL = (E_{TEL}^{(Cq_{TEL}[Calibrator] - Cq_{TEL}[Sample])}) / (E_{B2M}^{(Cq_{B2M}[Calibrator] - Cq_{B2M}[Sample])})$$

Where  $E_{TEL}$  and  $E_{B2M}$  are the mean reaction efficiencies for the respective amplicon group across all samples on a given plate;  $Cq_{TEL}[Calibrator]$  and  $Cq_{B2M}[Calibrator]$  are the average Cqs for the relevant amplicon across all calibrator samples on the plate; and  $Cq_{TEL}[Sample]$  and  $Cq_{B2M}[Sample]$  are the average of the triplicate Cqs for the sample for each amplicon.

### Data analysis

We began by checking the distribution of our telomere, FEC and leukocyte proportions (GLR, CD4:CD8 ratio, proportion of CD4 and CD8 T cells that were naïve) data. RTL was normally distributed but the other variables showed right skew. Log transformation yielded approximately normally distributions, and so log transformed FEC and leukocyte proportions were used in all analyses that followed. Eleven samples came from animals that were not caught at birth and had uncertain ages, so they were excluded from further analyses. We then calculated the Pearson's correlation coefficient among RTL and the leukocyte proportion measures. To test how RTL varied with age and

sex, we ran linear mixed effect models (LMMs) of RTL including individual identity (481 samples from 395 individuals) and qPCR plate (9 plates) as random effects and year, sex and age (as a linear and quadratic covariate) and the interaction between the age terms and sex as fixed effects. We assessed the significance of each term using likelihood ratio tests (LRTs). We subsequently tested whether independent effects of August weight and FEC on RTL were present, having accounted for age effects, by separately adding weight or FEC and their interactions with sex to the model and testing whether the addition improved model fit using LRTs. We further tested whether horn type was associated with RTL by adding a three way interaction among age, sex and horn type and all associated lower order interactions and sequentially deleting terms until only significant terms remained in the model.

To test our hypothesis that costs of investment in a key secondary sexual trait in males, horn length, might be reflected in RTL, we ran a separate set of LMMs restricting our data set to only normal horned males (N = 130). Only a small proportion of Soay ewes have normal horns and exhibit horn growth (74 out of 348 females in our data set) and this combined with the absence of evidence that horn growth is costly or important in reproduction lead us to restrict our analyses to male horn growth. As horns are grown incrementally each year in sheep, horn length is very strongly determined by an animal's age. To avoid the potentially confounding association between age and horn length in our models (as RTL is age-dependent in males, see Results), we ran a separate models for lambs (aged < 1 year) and adults (aged 1 year or more). In lambs, horn length reflects horn growth over the first four months of life, and we tested its significance by adding both a linear and quadratic horn length fixed effect to a LMM of RTL with plate as random effect (there was only one observation per individual in this data set), alongside year and lamb age in days (as not all lambs were exactly the same age when caught in August) as additional fixed effects. In adult males, horn length inevitably

increases with age as horn increments are grown. We tested the association between horn length and RTL in adults in a similar fashion to lambs, but included age in years as a fixed effect to account for the age-related change in both RTL and horn length.

To test for associations between RTL and leukocyte proportions and determine whether changes in leukocyte cell structure could explain observed patterns of variation in RTL with age and sex, we added all four leukocyte proportion measures into our final overall LMM of RTL and assessed whether previously observed sex and age effects remained significant in this model. We used a similar approach in our models including horn length restricted to normal horned males. For descriptive purposes, we calculated means and standard errors for all immune cell percentages and counts and ratios derived from them both overall and within each major age class (lamb, yearling, adult: 2-6 years, geriatric: >6 years). We then ran LMMs of each immune cell measurement including age, sex and their interaction and year as fixed effects and individual identity as a random effect, and simplified these models as described above. All analyses were conducted in R version 3.2.3 (R Core Team 2012).

## Results

There was a significant interaction between the effects of age and sex on RTL ( $\chi^2_{(d.f. = 1)} = 5.67$ ,  $P = 0.02$ ): males, but not females, showed a decline in telomere length with age (estimated difference in slope between males and females:  $b = -0.019 \pm 0.008$  SE; Figure 1). There was no evidence for significant quadratic effects of age (dropping  $age^2$  and  $age^2$ -by-sex interaction from the model:  $\chi^2_{(2)} = 1.78$ ,  $P = 0.41$ ) and no significant difference in RTL between capture years (dropping year:  $\chi^2_{(1)} = 1.58$ ,

P = 0.21). When we separated our RTL data set into age groups and re-ran our LMMs (without individual as a random effect, because there were no or very few repeat measures in each data subset), we found that male RTL was only significantly shorter than female RTL in adults aged three years or more (N = 193; males vs. females:  $b = -0.089 \pm 0.042$  SE,  $\chi^2_{(1)} = 4.58$ ,  $P = 0.03$ ). There was no significant sex difference among lambs (N = 200;  $b = 0.004 \pm 0.023$  SE,  $\chi^2_{(1)} = 0.32$ ,  $p = 0.86$ ) or individuals aged two years or less (N = 288;  $b = 0.016 \pm 0.020$  SE,  $\chi^2_{(1)} = 0.68$ ,  $P = 0.43$ ). August weight, FEC and horn type were not significantly associated with RTL independent of age and sex. The addition of weight and its interaction with sex to a model already including an interaction between age and sex did not significantly improve model fit ( $\chi^2_{(2)} = 1.29$ ,  $P = 0.53$ ), and nor did the addition of FEC and its interaction with sex ( $\chi^2_{(2)} = 0.36$ ,  $P = 0.84$ ). The addition of horn type and its potential interactions with age and sex also did not significantly improve model fit when added to the model including the interaction between age and sex ( $\chi^2_{(6)} = 3.12$ ,  $P = 0.79$ ).

Horn length was negatively associated with RTL in normal horned male lambs, but not in older males once their age was accounted for (Figure 2). In a model of RTL including only normal horned males, there was a significant decline in RTL with age ( $b = -0.023 \pm 0.008$  SE,  $\chi^2_{(1)} = 9.28$ ,  $P < 0.01$ ). In a model restricted to lambs only, there was no effect of the lamb's age in days at capture in August ( $\chi^2_{(1)} = 0.35$ ,  $P = 0.55$ ) and a quadratic effect of horn length was not significant ( $\chi^2_{(1)} = 0.62$ ,  $P = 0.43$ ). Horn length was significantly negatively associated with RTL (N = 88;  $b = -0.0010 \pm 0.0004$  SE,  $\chi^2_{(1)} = 5.43$ ,  $P < 0.05$ ; Figure 2A). This association remained significant, and actually became stronger, when lamb August weight was included in the model ( $b = -0.0018 \pm 0.0005$  SE,  $\chi^2_{(1)} = 11.56$ ,  $P < 0.001$ ). In models including adult normal horned males, horn length was significant when age was not included in the model (N = 42;  $b = -0.0007 \pm 0.0002$  SE,  $\chi^2_{(1)} = 10.06$ ,  $P < 0.01$ ; Figure 2B), but was non-



significant when the age was included in the model ( $b = -0.0002 \pm 0.0003$  SE,  $\chi^2_{(1)} = 0.59$ ,  $P = 0.44$ ; Figure 2C).

Means and standard errors for all immune cell measurements are presented in Table S2. All four measures of leukocyte proportions used in analyses (GLR, CD4:CD8 ratio and the proportion of naïve CD4 and CD8 T cells) were significantly associated with age: GLR increased with age and the other three measures declined (Figure 3, Table S3). GLR and the proportion of CD4+ naïve T cells showed significant age by sex interactions: males increased their GLR and decreased their proportion of CD4 naïve T cells more rapidly with age than females (Figure 3, Table S3). Variation in RTL and its association with sex and age were largely independent of variation in the proportions of different leukocyte cell types measured in the samples. Pearson's correlation coefficients among RTL and the ratios and proportions of leukocyte cell types ranged between +0.1 and -0.1 (Figure 4). In a LMM of RTL restricted to samples for which all leukocyte proportion measures were available ( $N = 437$ ), the age by sex interaction remained marginally non-significant (male vs. female slope:  $b = -0.016 \pm 0.009$  SE,  $\chi^2_{(1)} = 3.46$ ,  $P = 0.06$ ). Addition of all four leukocyte proportion measurements to the model of RTL did not improve model fit ( $\chi^2_{(4)} = 3.34$ ,  $P = 0.50$ ) or meaningfully alter the effect sizes or significance of the age by sex interaction ( $b = -0.015 \pm 0.009$  SE,  $\chi^2_{(1)} = 3.16$ ,  $P = 0.08$ ). The significant negative association between RTL and horn length in male lambs remained when all four leukocyte cell type measurements were included in the LMM ( $N = 80$ ;  $b = -0.0010 \pm 0.0005$ ,  $\chi^2_{(1)} = 4.23$ ,  $P < 0.05$ ) and, as before, the addition of the four leukocyte measurements to the model did not improve fit ( $\chi^2_{(4)} = 4.18$ ,  $P = 0.38$ ).

## Discussion

380

381 This study provides, to our knowledge, the first evidence for sex differences in telomere length from  
382 a wild mammal. Differences in LTL between males and females were not detectable before three  
383 years of age, suggesting that there was no sex difference in LTL at birth in our study system. Longer  
384 LTLs in females than males in adulthood but not early life have also been documented in humans and  
385 laboratory rodents, suggesting that these sex differences in LTL may arise as a result of differences in  
386 attrition rates through development and early adulthood (Cherif *et al.* 2003; Gardner *et al.* 2014;  
387 Lapham *et al.* 2015; Tarry-Adkins *et al.* 2006). In our study, the presence of shorter LTL in older males  
388 compared to females could be due to sex differences in telomere attrition rate or in selective  
389 mortality associated with telomere length. Sex differences in selection on erythrocyte telomere  
390 length have been documented in wild sand lizards (*Lacerta agilis*; Olsson *et al.* 2011), and winter  
391 mortality in Soay sheep on St Kilda is male-biased at all ages (Clutton-Brock & Pemberton 2004).  
392 Current evidence from wild vertebrates, including a previous study of Soay sheep, points to positive  
393 associations among LTL or ELT and either annual survival or longevity and thus selective  
394 disappearance of individuals with short telomeres (Barrett *et al.* 2013; Beirne *et al.* 2014; Bize *et al.*  
395 2009; Fairlie *et al.* 2016; Olsson *et al.* 2011; Salomons *et al.* 2009). Although sex differences in  
396 telomere attrition rate could explain our results, the presence of stronger selective disappearance of  
397 individuals with short telomeres in females than males could also be responsible. We had insufficient  
398 longitudinal repeat samples within our very largely cross-sectional data set to differentiate these two  
399 possibilities. Longitudinal telomere data from both sexes spanning the period from birth to later  
400 adulthood are required to fully understand the within- and among-individual processes responsible  
401 for sex differences in telomere length.

402

We found no association between LTL and weight or Strongyle FEC in either sex, suggesting sex differences in growth or body size and in infection with gastrointestinal parasites could not explain observed sex differences in LTL in later life. This adds to the general lack of support for the sexual size dimorphism hypothesis from both among- and within-species studies (Barrett & Richardson 2011; Beirne *et al.* 2014; Olsson *et al.* 2011). Previous studies have found associations between micro-parasite infection status and telomere length (Asghar *et al.* 2015; Asghar *et al.* 2016; Ilmonen *et al.* 2008), whilst our study measured burden with chronically infecting gastro-intestinal nematode parasites. It seems plausible that a larger longitudinal study may be required to detect the immune consequences of such highly localised and long lasting infections and telomere lengths. Studies in laboratory rats have suggested sex differences in TL emerge around puberty as a direct result of the differential effects of sex hormones on telomere dynamics (Cherif *et al.* 2003; Tarry-Adkins *et al.* 2006). Soay sheep are sexually mature in their first year, so our data imply that sex differences in LTL emerge several years after puberty, but could still be the result of cumulative telomere eroding effects of testosterone relative to oestrogen. Over and above hormonal causes, males surviving to later adulthood will have experienced the cumulative physiological demands associated with years of rutting which could generate further differences in the rate of telomere attrition compared to females.

We have presented rare evidence of reduced telomere length associated with investment in a reproductive trait under natural conditions. Studies of birds in both laboratory and wild populations have found that experimentally increasing reproductive effort decreased ETL in parents, in the short term at least (Heidinger *et al.* 2012; Reichert *et al.* 2014; Sudyka *et al.* 2014, but see Beaulieu *et al.* 2011; Voillemot *et al.* 2012), whilst a non-manipulative field study found negative associations between ETL and arrival date and the number of nestlings (Bauch *et al.* 2013). We found a significant

negative association between LTL and horn length in males at around four months of age, but not in adult males. This is consistent with LTL shortening reflecting some physiological cost of horn growth, but raises the question of why it was only detected in lambs. Lambs aged four months are growing rapidly and contending with their first exposures to the parasite fauna on St Kilda (Clutton-Brock & Pemberton 2004), and this may mean that the costs of investment in secondary sexual traits such as horn growth are most pronounced at this age. This stage also captures the relationship between LTL and horn growth prior to potentially confounding effects of over-winter viability selection and subsequent rutting effort and incremental horn growth, which could make the cost easier to detect. A study of wild sand lizards documented disruptive selection on ETL in females but not in males (Olsson *et al.* 2011), and previous work on Soay sheep showed that the alleles associated with horn growth improves breeding success at a cost to longevity in males (Johnston *et al.* 2013). These studies and our present results suggest that the degree to which sex differences in LTL are generated and maintained by sexual differences in selection on telomere length is an important area for future study.

Our study offers indirect evidence that the observed relationships among LTL, age and sex are not driven by variation in the proportions of particular leukocyte cell types, which are known to have different telomere lengths during adulthood in humans (Aubert *et al.* 2012; Kimura *et al.* 2010; Weng 2001). We documented age-related changes in the proportions of different leukocyte cell types that were consistent with observations in humans and laboratory mice, and in our own previous studies of this system (Linton & Dorshkind 2004; Nussey *et al.* 2012; Pawelec *et al.* 2010). Of particular note, the decline in the proportion of naïve helper and cytotoxic T cells (Figure 3) with age could generate population level declines in mean LTL because naïve T cells have longer telomeres than effector and memory T cells (Aubert *et al.* 2012; Weng 2001). Furthermore, the presence of a sex by age

interaction for CD4+ naïve cells, indicating that males have more rapid rates of declines with age than females (Figure 3, Table S3), could have been responsible for the observed sex difference in LTL with age. However, we found little evidence that LTL was correlated with any of the leukocyte proportion measurements and we showed that the main results of our LTL models were not influenced by the inclusion of the leukocyte proportion measurements. Our remote field study site precluded the use of more sophisticated methods to determine the actual telomere lengths of particular types of leukocytes, such as cell sorting or flow-FISH (Kimura *et al.* 2010; Aubert *et al.* 2016). However, our findings are consistent with the mounting evidence that TL is highly correlated across leukocyte cell subsets, and across tissues more generally, within individual organisms (Asghar *et al.* 2016; Aubert *et al.* 2012; Daniali *et al.* 2013; Kimura *et al.* 2010; Reichert *et al.* 2013). This suggests that telomere length, whether measured in leukocytes or erythrocytes, could reflect variation that exists in the individual's haematopoietic stem cell compartment, and even more general organism-wide variation in telomere length (Daniali *et al.* 2013; Kimura *et al.* 2010; Reichert *et al.* 2013).

**Acknowledgements:** We are grateful to the National Trust for Scotland for permission to work on St Kilda; QinetiQ and Kilda Cruises for logistical support; the 2014 and 2015 August catch teams for all their hard work in the field; Rose Zamoyska and Martin Waterfall for their help with the immunological assays; and Adam Hayward for comments on a draft of the manuscript. RLW was supported by a BBSRC EASTBIO PhD studentship and DHN by a BBSRC David Phillip's fellowship. The laboratory work was funded by a BBSRC grant to DHN (BB/L20769/1) and the Soay sheep project was funded by consecutive NERC grants to JMP and others.

## 475    **References**

- 476    Akkad A, Hastings R, Konje J, *et al.* (2006) Telomere length in small-for-gestational-age babies. *BJOG: An International Journal of Obstetrics & Gynaecology* **113**, 318-323.
- 477    Andersson M (1994) *Sexual Selection*. Princeton University Press, Princeton.
- 478    Armanios M, Blackburn EH (2012) The telomere syndromes. *Nature Reviews Genetics* **13**, 693-704.
- 479    Asghar M, Hasselquist D, Hansson B, *et al.* (2015) Hidden costs of infection: chronic malaria  
480    accelerates telomere degradation and senescence in wild birds. *Science* **347**, 436-438.
- 481    Asghar M, Palinauskas V, Zaghoudi-Allan Ng, *et al.* (2016) Parallel telomere shortening in multiple  
482    body tissues owing to malaria infection. *Proceedings of the Royal Society London, Series B*  
483    **283**, 20161184.
- 484    Aubert G, Baerlocher GM, Vulto I, Poon SS, Lansdorp PM (2012) Collapse of telomere homeostasis in  
485    hematopoietic cells caused by heterozygous mutations in telomerase genes. *PLoS Genetics* **8**,  
486    e1002696.
- 487    Aviv A, Shay J, Christensen K, Wright W (2005) The longevity gender gap: are telomeres the  
488    explanation? *Science's SAGE KE* **2005**, pe16.
- 489    Baerlocher GM, Rice K, Vulto I, Lansdorp PM (2007) Longitudinal data on telomere length in  
490    leukocytes from newborn baboons support a marked drop in stem cell turnover around 1  
491    year of age. *Aging Cell* **6**, 121-123.
- 492    Bain BJ (2014) *Blood cells: a practical guide* John Wiley & Sons.
- 493    Barrett ELB, Burke TA, Hammers M, Komdeur J, Richardson DS (2013) Telomere length and dynamics  
494    predict mortality in a wild longitudinal study. *Molecular Ecology* **22**, 249-259.
- 495    Barrett ELB, Richardson DS (2011) Sex differences in telomeres and lifespan. *Aging Cell* **10**, 913-921.
- 496    Bauch C, Becker PH, Verhulst S (2013) Telomere length reflects phenotypic quality and costs of  
497    reproduction in a long-lived seabird. *Proceedings of the Royal Society London B* **280**,  
498    20122540.
- 499    Beaulieu M, Reichert S, Le Maho Y, Ancel A, Criscuolo F (2011) Oxidative status and telomere length  
500    in a long-lived bird facing a costly reproductive event. *Functional Ecology* **25**, 577-585.
- 501    Beirne C, Delahay R, Hares M, Young A (2014) Age-Related Declines and Disease-Associated Variation  
502    in Immune Cell Telomere Length in a Wild Mammal. *PLoS ONE* **9**, e108964.
- 503    Berglund K, Reynolds CA, Ploner A, *et al.* (2016) Longitudinal decline of leukocyte telomere length in  
504    old age and the association with sex and genetic risk. *Aging* **8**, 7.
- 505    Bize P, Criscuolo F, Metcalfe NB, Nasir L, Monaghan P (2009) Telomere dynamics rather than age  
506    predict life expectancy in the wild. *Proceedings Of The Royal Society London B* **276**, 1679-  
507    1683.
- 508    Blackburn EH (1991) Structure and function of telomeres. *Nature* **350**, 569-573.
- 509    Blackburn EH, Epel ES, Lin J (2015) Human telomere biology: A contributory and interactive factor in  
510    aging, disease risks, and protection. *Science* **350**, 1193-1198.
- 511    Cawthon RM (2002) Telomere measurement by quantitative PCR. *Nucleic Acids Research* **30**, e47.
- 512    Cawthon RM, Smith KR, O'Brien E, Sivatchenko A, Kerber RA (2003) Association between telomere  
513    length in blood and mortality in people aged 60 years or older. *The Lancet* **361**, 393-395.
- 514    Cherif H, Tarry J, Ozanne S, Hales C (2003) Ageing and telomeres: A study into organ- and gender-  
515    specific telomere shortening. *Nucleic Acids Research* **31**, 1576-1583.
- 516    Clutton-Brock TH, Pemberton JM (2004) *Soay sheep: Dynamics and selection in an island population*.  
517    Cambridge University Press, Cambridge.
- 518    Coviello-McLaughlin GM, Prowse KR (1997) Telomere length regulation during postnatal  
519    development and ageing in *Mus spretus*. *Nucleic Acids Research* **25**, 3051-3058.
- 520

521 Daniali L, Benetos A, Susser E, *et al.* (2013) Telomeres shorten at equivalent rates in somatic tissues  
 522 of adults. *Nature communications* **4**, 1597.  
 523 De Lange T (2004) T-loops and the origin of telomeres. *Nature Reviews Molecular Cell Biology* **5**, 323-  
 524 329.  
 525 Epel ES, Blackburn EH, Lin J, *et al.* (2004) Accelerated telomere shortening in response to life stress.  
 526 *Proceedings of the National Academy of Sciences of the United States of America* **101**, 17312-  
 527 17315.  
 528 Factor-Litvak P, Susser E, Kezios K, *et al.* (2016) Leukocyte Telomere Length in Newborns: Implications  
 529 for the Role of Telomeres in Human Disease. *Pediatrics* **137**.  
 530 Fairlie J, Holland R, Pilkington JG, *et al.* (2016) Life-long leukocyte telomere dynamics and survival in a  
 531 free-living mammal. *Aging Cell* **15**, 140-148.  
 532 Gardner JP, Kimura M, Chai W, *et al.* (2007) Telomere dynamics in Macaques and humans. *Journals*  
 533 *Of Gerontology Series A-Biological Sciences And Medical Sciences* **62**, 367-374.  
 534 Gardner M, Bann D, Wiley L, *et al.* (2014) Gender and telomere length: Systematic review and meta-  
 535 analysis. *Experimental Gerontology* **51**, 15-27.  
 536 Giefing-Kröll C, Berger P, Lepperdinger G, Grubeck-Loebenstien B (2015) How sex and age affect  
 537 immune responses, susceptibility to infections, and response to vaccination. *Aging Cell* **14**,  
 538 309-321.  
 539 Gomes NMV, Ryder OA, Houck ML, *et al.* (2011) Comparative biology of mammalian telomeres:  
 540 hypotheses on ancestral states and the roles of telomeres in longevity determination. *Aging*  
 541 *Cell* **10**, 761-768.  
 542 Gulland FMD, Fox M (1992) Epidemiology of Nematode Infections of Soay Sheep (*Ovis aries* L) on St-  
 543 Kilda. *Parasitology* **105**, 481-492.  
 544 Hayward AD, Wilson AJ, Pilkington JG, Pemberton JM, Kruuk LEB (2009) Ageing in a variable habitat:  
 545 environmental stress affects senescence in parasite resistance in St Kilda Soay sheep.  
 546 *Proceedings of the Royal Society of London B* **276**, 3477-3485.  
 547 Heidinger BJ, Blount JD, Boner W, *et al.* (2012) Telomere length in early life predicts lifespan.  
 548 *Proceedings of the National Academy of Sciences of the United States of America* **109**, 1743-  
 549 1748.  
 550 Hemann MT, Strong MA, Hao L-Y, Greider CW (2001) The shortest telomere, not average telomere  
 551 length, is critical for cell viability and chromosome stability. *Cell* **107**, 67-77.  
 552 Ilmonen P, Kotrschal A, Penn DJ (2008) Telomere attrition due to infection. *PLoS ONE* **3**, e2143.  
 553 Johnston SE, Gratten J, Berenos C, *et al.* (2013) Life history trade-offs at a single locus maintain  
 554 sexually selected genetic variation. *Nature* **502**, 93-95.  
 555 Kimura M, Gazitt Y, Cao X, *et al.* (2010) Synchrony of telomere length among hematopoietic cells.  
 556 *Experimental Hematology* **38**, 854-859.  
 557 Lapham K, Kvale MN, Lin J, *et al.* (2015) Automated assay of telomere length measurement and  
 558 informatics for 100,000 subjects in the Genetic Epidemiology Research on Adult Health and  
 559 Aging (GERA) cohort. *Genetics* **200**, 1061-1072.  
 560 Linton PJ, Dorshkind K (2004) Age-related changes in lymphocyte development and function. *Nature*  
 561 *Immunology* **5**, 133-139.  
 562 Monaghan P (2010) Telomeres and life histories: the long and the short of it. *Annals of the New York*  
 563 *Academy of Science* **1206**, 130-142.  
 564 Monaghan P (2014) Organismal stress, telomeres and life histories. *Journal of Experimental Biology*  
 565 **217**, 57-66.  
 566 Monaghan P, Haussmann MF (2006) Do telomere dynamics link lifestyle and lifespan? *Trends In*  
 567 *Ecology & Evolution* **21**, 47-53.

- Moore SL, Wilson K (2002) Parasites as a viability cost of sexual selection in natural populations of mammals. *Science* **297**, 2015-2018.
- Nussey DH, Watt K, Pilkington JG, Zamoyska R, McNeilly TN (2012) Age-related variation in immunity in a wild mammal population. *Aging Cell* **11**, 178-180.
- Okuda K, Bardeguet A, Gardner JP, et al. (2002) Telomere length in the newborn. *Pediatric research* **52**, 377-381.
- Olsson M, Pauliny A, Wapstra E, et al. (2011) Sexual differences in telomere selection in the wild. *Molecular Ecology* **20**, 2085-2099.
- Pawelec G, Larbi A, Derhovanessian E (2010) Senescence of the Human Immune System. *Journal of Comparative Pathology* **142**, S39-S44.
- Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research* **29**.
- Reichert S, Criscuolo F, Verinaud E, Zahn S, Massemin S (2013) Telomere length correlations among somatic tissues in adult zebra finches. *PLoS ONE* **8**, e81496.
- Reichert S, Stier A, Zahn S, et al. (2014) Increased brood size leads to persistent eroded telomeres. *Frontiers in Ecology and Evolution* **2**, 9.
- Rode L, Nordestgaard BG, Bojesen SE (2015) Peripheral blood leukocyte telomere length and mortality among 64 637 individuals from the general population. *Journal of the National Cancer Institute* **107**, djv074.
- Ruijter JM, Ramakers C, Hoogaars WMH, et al. (2009) Amplification efficiency: linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Research* **37**.
- Salomons HM, Mulder GA, van de Zande L, et al. (2009) Telomere shortening and survival in free-living corvids. *Proceedings Of The Royal Society London B* **276**, 3157-3165.
- Seeker LA, Holland R, Underwood S, et al. (2016) Method specific calibration corrects for DNA extraction method effects on relative telomere length measurements by quantitative PCR. *PLoS ONE*, **11**, e0164046.
- Shalev I (2012) Early life stress and telomere length: Investigating the connection and possible mechanisms. *Bioessays* **34**, 943-952.
- Shi-Ni L, Yahya Z, Zeegers D, et al. (2013) Distribution of telomere length in the cord blood of Chinese newborns. *British Journal of Medicine and Medical Research* **3**, 1004.
- Smith DL, Mattison JA, Desmond RA, et al. (2011) Telomere Dynamics in Rhesus Monkeys: No Apparent Effect of Caloric Restriction. *Journals Of Gerontology Series A-Biological Sciences And Medical Sciences* **66**, 1163-1168.
- Stindl R (2004) Tying it all together: Telomeres, sexual size dimorphism and the gender gap in life expectancy. *Medical Hypotheses* **62**, 151-154.
- Sudyka J, Arct A, Drobniak S, et al. (2014) Experimentally increased reproductive effort alters telomere length in the blue tit (*Cyanistes caeruleus*). *Journal of Evolutionary Biology* **27**, 2258-2264.
- Tarry-Adkins JL, Ozanne SE, Norden A, Cherif H, Hales CN (2006) Lower antioxidant capacity and elevated p53 and p21 may be a link between gender disparity in renal telomere shortening, albuminuria, and longevity. *American Journal of Physiology-Renal Physiology* **290**, F509-F516.
- R Core Development Team (2012) R: A language and environment for statistical computing. . R Foundation for Statistical Computing, Vienna, Austria.
- Voillemot M, Hine K, Zahn S, et al. (2012) Effects of brood size manipulation and common origin on phenotype and telomere length in nestling collared flycatchers. *BMC Ecology* **12**.
- von Zglinicki T (2002) Oxidative stress shortens telomeres. *Trends in Biochemical Sciences* **27**, 339-344.



Watson, R.L., McNeilly, T.N., Watt, K.A., Pemberton, J.M., Pilkington, J.G., Waterfall, M., Hopper, P.R.T., Cooney, D., Zamoyska, R. & Nussey, D.H. (2016) Cellular and humoral immunity in a wild mammal: Variation with age & sex and association with over-winter survival. *Ecology & Evolution*, online early.


Weng N-p (2001) Interplay between telomere length and telomerase in human leukocyte differentiation and aging. *Journal of Leukocyte Biology* **70**, 861-867.

Wilson K, Grenfell BT, Pilkington JG, Boyd HEG, Gulland FM (2004) Parasites and their impact. In: *Soay Sheep: Dynamics and Selection in an Island Population* (eds. Clutton-Brock TH, Pemberton JM), pp. 113-165. Cambridge University Press, Cambridge.


**Data accessibility:** Data used in this paper is available on Dryad:  
<http://dx.doi.org/10.5061/dryad.kd92s>.

**Authors' contributions:** All authors contributed to the design of the study; RLW and DHN wrote the manuscript with editorial input from all co-authors; RLW, HF and DHN conducted the statistical analyses; RLW, EJB, JF, SU, KW, ES-C and RVA conducted the laboratory work; JMP and JGP coordinated field trips and conducted data and sample collection in the field.


**Figure 1.** Older males have shorter relative leukocyte telomere length (RTL) than females, but no sex differences are apparent amongst young animals. Raw data for males (blue triangles) and females (red circles) are presented with a linear regression (blue and red lines respectively) and associated standard error (grey shading) for each sex.



**Figure 2.** Relative leukocyte telomere length (RTL) is negatively associated with horn length amongst normal horned males in lambs (<1 year old), but not adults (1 year old or greater) once effects of age are accounted for. Plots show raw RTL against horn length measures with linear regression (black line) and associated standard error (grey shading) for: (A) male lamb, (B) adult males, (C) adult males, having corrected horn length for age by taking residuals from a regression of RTL on age.



**Figure 3.** Scatter plots showing relationships with age and sex for (A): granulocyte to lymphocyte ratios, (B): CD4 to CD8 T cell ratios, (C) the proportion of CD4+ “helper” T cells that were naïve and (D) the proportion of CD8+ “cytotoxic” T cells that were naïve. All proportions and ratios are log transformed. Raw data for males (blue triangles) and females (red circles) are presented with a linear regression (blue and red lines) and associated standard error (grey shading) for each sex.



**Figure 4.** Scatter plots illustrating correlations among relative leukocyte telomere length (RTL) and measures of the proportion of different leukocyte cell types. Scatterplots show relationships between RTL and: (A) granulocyte to lymphocyte ratio ("G:L ratio"), (B) CD4+ T cell to CD8+ T cell ratio ("CD4:CD8 ratio"), (C) proportion of CD4+ T cells positive for CD45RA ("CD4 naïve") (D) proportion of CD8+ T cells positive for CD45RA ("CD8 naïve"). All leukocyte ratios and proportions were log transformed; Pearson's correlation coefficients and associated P values presented for each plot.

